

EQUIVALENCE OF DISTANCE-BASED AND RKHS-BASED STATISTICS IN HYPOTHESIS TESTING

BY DINO SEJDINOVIC^{*}, BHARATH SRIPERUMBUDUR^{*,§},
ARTHUR GRETTON^{*,†,§} AND KENJI FUKUMIZU[†]

*Gatsby Unit, CSML, University College London^{*},
The Institute of Statistical Mathematics, Tokyo[†], and
Max Planck Institute for Intelligent Systems, Tübingen[‡]*

Abstract We provide a unifying framework linking two classes of statistics used in two-sample and independence testing: on the one hand, the energy distances and distance covariances from the statistics literature; on the other, Maximum Mean Discrepancies (MMD), i.e., distances between embeddings of distributions to reproducing kernel Hilbert spaces (RKHS), as established in machine learning. In the case where the energy distance is computed with the semimetric of negative type, a positive definite kernel, termed distance kernel, may be defined such that the MMD corresponds exactly to the energy distance. Conversely, for any positive definite kernel, we can interpret the MMD as energy distance with respect to some negative-type semimetric. This equivalence readily extends to distance covariance using kernels on the product space. We determine the class of probability distributions for which the test statistics are consistent against all alternatives. Finally, we investigate the performance of the family of distance kernels in two-sample and independence tests: we show in particular that the energy distance most commonly employed in statistics is just one member of a parametric family of kernels, and that other choices from this family can yield more powerful tests.

1. Introduction. The problem of testing statistical hypotheses in high dimensional spaces is particularly challenging, and has been a recent focus of considerable work in both the statistics and the machine learning communities. On the statistical side, two-sample testing in Euclidean spaces (of whether two independent samples are from the same distribution, or from different distributions) can be accomplished using a so-called energy distance as a statistic [Székely and Rizzo, 2004, 2005]. Such tests are consistent against all alternatives as long as the random variables have finite first moments. A related dependence measure between vectors of high dimension is

[§]These authors contributed equally.

AMS 2000 subject classifications: Primary 62G10, 62H20, 68Q32; secondary 46E22

Keywords and phrases: reproducing kernel Hilbert spaces, distance covariance, two-sample testing, independence testing

the distance covariance [Székely et al., 2007, Székely and Rizzo, 2009], and the resulting test is again consistent for variables with bounded first moment. The distance covariance has had a major impact in the statistics community, with [Székely and Rizzo, 2009] being accompanied by an editorial introduction and discussion. A particular advantage of energy distance-based statistics is their compact representation in terms of certain expectations of pairwise Euclidean distances, which leads to straightforward empirical estimates. As a follow-up work, Lyons [2011] generalized the notion of distance covariance to metric spaces of negative type (of which Euclidean spaces are a special case).

On the machine learning side, two-sample tests have been formulated based on embeddings of probability distributions into reproducing kernel Hilbert spaces [Gretton et al., 2012], using as the test statistic the difference between these embeddings: this statistic is called the Maximum Mean Discrepancy (MMD). This distance measure was also applied to the problem of testing for independence, with the associated test statistic being the Hilbert-Schmidt Independence Criterion (HSIC) [Gretton et al., 2005, 2008, Smola et al., 2007, Zhang et al., 2011]. Both tests are shown to be consistent against all alternatives when a characteristic RKHS is used [Fukumizu et al., 2009, Sriperumbudur et al., 2010]. Such tests can further be generalized to structured and non-Euclidean domains, such as text strings, graphs, and groups [Fukumizu et al., 2009].

Despite their striking similarity, the link between energy distance-based tests and kernel-based tests has been an open question. In the discussion of [Székely and Rizzo, 2009], Gretton et al. [2009b, p. 1289] first explored this link in the context of independence testing, and found that interpreting the distance-based independence statistic as a kernel statistic is not straightforward, since Bochner’s theorem does not apply to the choice of weight function used in the definition of the distance covariance (we briefly review this argument in Section 4.3). Székely and Rizzo [2009, Rejoinder, p. 1303] confirmed this conclusion, and commented that RKHS-based dependence measures do not seem to be formal extensions of the distance covariance because the weight function is not integrable. Our contribution resolves this question and shows that RKHS-based dependence measures are precisely the formal extensions of the distance covariance, where the problem of non-integrability of weight functions is circumvented by using translation-variant kernels, i.e., *distance-induced kernels*, a novel family of kernels that we introduce in Section 2.4.

In the case of two-sample testing, we demonstrate that energy distances are in fact maximum mean discrepancies arising from the same family of

distance-induced kernels. A number of interesting consequences arise from this insight: first, as the energy distance (and distance covariance) derives from a particular choice of a kernel, we can consider analogous quantities arising from other kernels, and yielding a more sensitive test. Second, results from [Gretton et al., 2009a, Zhang et al., 2011] may be applied to get consistent two-sample and independence tests for the energy distance, without using bootstrap, which perform much better than the upper bound proposed by Székely et al. [2007] as an alternative to the bootstrap. Third, in relation to Lyons [2011], we obtain a new family of characteristic kernels arising from semimetric spaces of negative type (where the triangle inequality need not hold), which are quite unlike the characteristic kernels defined via Bochner’s theorem [Sriperumbudur et al., 2010].

The structure of the paper is as follows: in Section 2, we provide the necessary definitions from RKHS theory, and the relation between RKHS and semimetrics of negative type. We also give a review of the principles behind the Maximum Mean Discrepancy (MMD) and the Hilbert-Schmidt Independence Criterion (HSIC), the RKHS-based statistics used for two-sample and independence testing, respectively. In Sections 3 and 4, we review the notions of energy distance and distance covariance in general (semi)metric spaces of negative type, and show them to be special cases of MMD and HSIC, respectively. We give conditions for these quantities to distinguish between probability measures in Section 5, thus obtaining a new family of characteristic kernels. Empirical estimates of these quantities and associated two-sample and independence tests are described in Section 6. Finally, in Section 7, we investigate the performance of the test statistics on a variety of testing problems, which demonstrate the strengths of the new kernel family.

This paper extends the conference publication [Sejdinovic et al., 2012], and gives a detailed technical discussion and proofs which were omitted in that work.

2. Kernels and (semi)metrics. In this section, we introduce concepts and notation required to understand reproducing kernel Hilbert spaces (Section 2.1), and distribution embeddings into RKHS. We then introduce semimetrics of negative type (Section 2.3), and reveal their relation to RKHS kernels.

2.1. RKHS and kernel embeddings. Unless stated otherwise, we will assume that $\mathcal{Z}$ is any topological space. We will denote by $\mathcal{M}(\mathcal{Z})$ the set of all finite signed Borel measures on $\mathcal{Z}$, and by $\mathcal{M}_+^1(\mathcal{Z})$ the set of all Borel probability measures on $\mathcal{Z}$.

DEFINITION 1. (**RKHS**) Let $\mathcal{H}$ be a Hilbert space of real-valued functions defined on $\mathcal{Z}$. A function $k : \mathcal{Z} \times \mathcal{Z} \rightarrow \mathbb{R}$ is called a *reproducing kernel* of $\mathcal{H}$ if:

1. $\forall z \in \mathcal{Z}, k(\cdot, z) \in \mathcal{H}$, and
2. $\forall z \in \mathcal{Z}, \forall f \in \mathcal{H}, \langle f, k(\cdot, z) \rangle_{\mathcal{H}} = f(z)$.

If $\mathcal{H}$ has a reproducing kernel, it is said to be a *reproducing kernel Hilbert space* (RKHS).

According to the Moore-Aronszajn theorem [Berlinet and Thomas-Agnan, 2004, p. 19], for every symmetric, positive definite function (henceforth *kernel*) $k : \mathcal{Z} \times \mathcal{Z} \rightarrow \mathbb{R}$, there is an associated RKHS $\mathcal{H}_k$ of real-valued functions on $\mathcal{Z}$ with reproducing kernel k . The map $\varphi : \mathcal{Z} \rightarrow \mathcal{H}_k, \varphi : z \mapsto k(\cdot, z)$ is called the canonical feature map or the Aronszajn map of k . We will say that k is a *nondegenerate kernel* if its Aronszajn map is injective. The notion of feature map can be extended to kernel embeddings of finite signed Borel measures on $\mathcal{Z}$ [Smola et al., 2007, Sriperumbudur et al., 2010], [Berlinet and Thomas-Agnan, 2004, Chapter 4].

DEFINITION 2. (**Kernel embedding**) Let k be a kernel on $\mathcal{Z}$, and $\nu \in \mathcal{M}(\mathcal{Z})$. The *kernel embedding* of ν into the RKHS $\mathcal{H}_k$ is $\mu_k(\nu) \in \mathcal{H}_k$ such that $\int f(z) d\nu(z) = \langle f, \mu_k(\nu) \rangle_{\mathcal{H}_k}$ for all $f \in \mathcal{H}_k$.

Alternatively, the kernel embedding can be defined by the Bochner integral $\mu_k(\nu) = \int k(\cdot, z) d\nu(z)$. If a measurable kernel k is a bounded function, $\mu_k(\nu)$ exists for all $\nu \in \mathcal{M}(\mathcal{Z})$. On the other hand, if k is not bounded, there will always exist $\nu \in \mathcal{M}(\mathcal{Z})$, for which $\int k(\cdot, z) d\nu(z)$ diverges. The kernels we will consider in this paper will be continuous, and hence measurable, but unbounded, so kernel embeddings will not be defined for some finite signed measures. Thus, we need to restrict the attention to a particular class of measures for which kernel embeddings exist (this will be later shown to reflect the condition that random variables considered in distance covariance tests must have finite moments). Let k be a measurable kernel on $\mathcal{Z}$, and denote, for $\theta > 0$,

$$(2.1) \quad \mathcal{M}_k^\theta(\mathcal{Z}) = \left\{ \nu \in \mathcal{M}(\mathcal{Z}) : \int k^\theta(z, z) d|\nu|(z) < \infty \right\}.$$

Clearly,

$$(2.2) \quad \theta_1 \leq \theta_2 \implies \mathcal{M}_k^{\theta_2}(\mathcal{Z}) \subseteq \mathcal{M}_k^{\theta_1}(\mathcal{Z}).$$

Note that the kernel embedding $\mu_k(\nu)$ is well defined $\forall \nu \in \mathcal{M}_k^{1/2}(\mathcal{Z})$, by the Riesz representation theorem. Indeed, $T_k : f \mapsto \int f(z) d\nu(z)$ is then a bounded linear functional on $\mathcal{H}_k$, which can be seen using the reproducing property and the Cauchy-Schwarz inequality,

$$T_k f = \int f(z) d\nu(z) = \int \langle f, k(\cdot, z) \rangle_{\mathcal{H}_k} d\nu(z) \leq \|f\|_{\mathcal{H}_k} \int k^{1/2}(z, z) d|\nu|(z).$$

Thus, kernel embeddings of Borel probability measures in the space $\mathcal{M}_+^1(\mathcal{Z}) \cap \mathcal{M}_k^{1/2}(\mathcal{Z})$ do exist, and we introduce the notion of distance between Borel probability measures in this space using the Hilbert space distance between their embeddings.

DEFINITION 3. (Maximum mean discrepancy) Let k be a kernel on $\mathcal{Z}$, and let $P, Q \in \mathcal{M}_+^1(\mathcal{Z}) \cap \mathcal{M}_k^{1/2}(\mathcal{Z})$. The *maximum mean discrepancy (MMD)* γ_k between P and Q is given by [Gretton et al. \[2012, Lemma 4\]](#),

$$\gamma_k(P, Q) = \|\mu_k(P) - \mu_k(Q)\|_{\mathcal{H}_k}.$$

The following alternative representation of the squared MMD [from [Gretton et al., 2012, Lemma 6](#)] will be useful

$$\begin{aligned} \gamma_k^2(P, Q) &= \mathbb{E}_{ZZ'} k(Z, Z') + \mathbb{E}_{WW'} k(W, W') - 2\mathbb{E}_{ZW} k(Z, W) \\ (2.3) \quad &= \int \int k d([P - Q] \times [P - Q]), \end{aligned}$$

where $Z, Z' \stackrel{i.i.d.}{\sim} P$ and $W, W' \stackrel{i.i.d.}{\sim} Q$. If the restriction of μ_k to some $\mathcal{P}(\mathcal{Z}) \subseteq \mathcal{M}_+^1(\mathcal{Z})$ is well defined and injective, then k is said to be characteristic to $\mathcal{P}(\mathcal{Z})$, and it is said to be characteristic (without further qualification) if it is characteristic to $\mathcal{M}_+^1(\mathcal{Z})$. When k is characteristic, γ_k is a metric on the entire $\mathcal{M}_+^1(\mathcal{Z})$, i.e., $\gamma_k(P, Q) = 0$ iff $P = Q$, $\forall P, Q \in \mathcal{M}_+^1(\mathcal{Z})$. Conditions under which kernels are characteristic have been studied in [[Sriperumbudur et al., 2008](#), [Fukumizu et al., 2009](#), [Sriperumbudur et al., 2010](#)]. An alternative interpretation of (2.3) is as an integral probability metric [[Müller, 1997](#)],

$$(2.4) \quad \gamma_k(P, Q) = \sup_{f \in \mathcal{H}_k, \|f\|_{\mathcal{H}_k} \leq 1} [\mathbb{E}_{Z \sim P} f(Z) - \mathbb{E}_{W \sim Q} f(W)].$$

See [[Gretton et al., 2012](#)] for details.

2.2. *Hilbert Schmidt Independence Criterion (HSIC)*. The MMD can be employed to measure statistical dependence between random variables [Gretton et al., 2005, 2008, Smola et al., 2007, Zhang et al., 2011]. Let $\mathcal{X}$ and $\mathcal{Y}$ be two non-empty topological spaces and let $k_{\mathcal{X}}$ and $k_{\mathcal{Y}}$ be kernels on $\mathcal{X}$ and $\mathcal{Y}$, with respective RKHSs $\mathcal{H}_{k_{\mathcal{X}}}$ and $\mathcal{H}_{k_{\mathcal{Y}}}$. Then, by applying [Steinwart and Christmann, 2008, Lemma 4.6, p. 114],

$$(2.5) \quad k((x, y), (x', y')) = k_{\mathcal{X}}(x, x')k_{\mathcal{Y}}(y, y')$$

is a kernel on the product space $\mathcal{X} \times \mathcal{Y}$ with RKHS $\mathcal{H}_k$ isometrically isomorphic to the tensor product $\mathcal{H}_{k_{\mathcal{X}}} \otimes \mathcal{H}_{k_{\mathcal{Y}}}$.

DEFINITION 4. Let $X \sim P_X$ and $Y \sim P_Y$ be random variables on $\mathcal{X}$ and $\mathcal{Y}$, respectively, having joint distribution P_{XY} . Furthermore, let k be a kernel on $\mathcal{X} \times \mathcal{Y}$, given in (2.5). The Hilbert-Schmidt Independence Criterion (HSIC) of X and Y is the MMD γ_k between the joint distribution P_{XY} and the product of its marginals $P_X P_Y$.

Following [Smola et al., 2007, Section 2.3], we can expand HSIC as:

$$(2.6) \quad \begin{aligned} \gamma_k^2(P_{XY}, P_X P_Y) &= \|\mu_k(P_{XY}) - \mu_k(P_X P_Y)\|_{\mathcal{H}_k}^2 \\ &= \|\mathbb{E}_{XY}[k_{\mathcal{X}}(\cdot, X) \otimes k_{\mathcal{Y}}(\cdot, Y)] - \mathbb{E}_X k_{\mathcal{X}}(\cdot, X) \otimes \mathbb{E}_Y k_{\mathcal{Y}}(\cdot, Y)\|_{\mathcal{H}_{k_{\mathcal{X}}} \otimes \mathcal{H}_{k_{\mathcal{Y}}}}^2 \\ &= \mathbb{E}_{XY} \mathbb{E}_{X'Y'} k_{\mathcal{X}}(X, X') k_{\mathcal{Y}}(Y, Y') + \mathbb{E}_X \mathbb{E}_{X'} k_{\mathcal{X}}(X, X') \mathbb{E}_Y \mathbb{E}_{Y'} k_{\mathcal{Y}}(Y, Y') \\ &\quad - 2 \mathbb{E}_{X'Y'} [\mathbb{E}_X k_{\mathcal{X}}(X, X') \mathbb{E}_Y k_{\mathcal{Y}}(Y, Y')] \end{aligned}$$

where in the last step we used that:

$$\langle f \otimes g, f' \otimes g' \rangle_{\mathcal{H}_{k_{\mathcal{X}}} \otimes \mathcal{H}_{k_{\mathcal{Y}}}} = \langle f, f' \rangle_{\mathcal{H}_{k_{\mathcal{X}}}} \langle g, g' \rangle_{\mathcal{H}_{k_{\mathcal{Y}}}}.$$

It can be shown that this quantity is equal to the squared Hilbert-Schmidt norm of the covariance operator between RKHSs [Gretton et al., 2005]. We claim that $\gamma_k^2(P_{XY}, P_X P_Y)$ is well defined as long as $P_X \in \mathcal{M}_{k_{\mathcal{X}}}^1(\mathcal{X})$ and $P_Y \in \mathcal{M}_{k_{\mathcal{Y}}}^1(\mathcal{Y})$. Indeed, the embedding $\mu_k(P_X P_Y)$ of the product of marginals can be identified with the tensor product $\mu_{k_{\mathcal{X}}}(P_X) \otimes \mu_{k_{\mathcal{Y}}}(P_Y)$, where $\mu_{k_{\mathcal{X}}}(P_X)$ exists since $P_X \in \mathcal{M}_{k_{\mathcal{X}}}^1(\mathcal{X}) \subset \mathcal{M}_{k_{\mathcal{X}}}^{1/2}(\mathcal{X})$, and $\mu_{k_{\mathcal{Y}}}(P_Y)$ exists since $P_Y \in \mathcal{M}_{k_{\mathcal{Y}}}^1(\mathcal{Y}) \subset \mathcal{M}_{k_{\mathcal{Y}}}^{1/2}(\mathcal{Y})$. Furthermore, $\mu_k(P_{XY})$ exists since $P_{XY} \in \mathcal{M}_k^{1/2}(\mathcal{X} \times \mathcal{Y})$, which can be seen from the Cauchy-Schwarz inequality,

$$\begin{aligned}
\int k^{1/2}((x, y), (x, y)) dP_{XY}(x, y) &= \int k_{\mathcal{X}}^{1/2}(x, x) k_{\mathcal{Y}}^{1/2}(y, y) dP_{XY}(x, y) \\
&\leq \left(\int k_{\mathcal{X}}(x, x) dP_X(x) \int k_{\mathcal{Y}}(y, y) dP_Y(y) \right)^{1/2}.
\end{aligned}$$

2.3. Semimetrics of negative type. We will work with the notion of semimetric of negative type on a non-empty set $\mathcal{Z}$, where the “distance” function need not satisfy the triangle inequality. Note that this notion of semimetric is different to that which arises from the seminorm (also called the pseudonorm), where the distance between two distinct points can be zero.

DEFINITION 5. (Semimetric) Let $\mathcal{Z}$ be a non-empty set and let $\rho : \mathcal{Z} \times \mathcal{Z} \rightarrow [0, \infty)$ be a function such that $\forall z, z' \in \mathcal{Z}$,

1. $\rho(z, z') = 0$ if and only if $z = z'$, and
2. $\rho(z, z') = \rho(z', z)$.

Then $(\mathcal{Z}, \rho)$ is said to be a semimetric space and ρ is called a semimetric on $\mathcal{Z}$. If, in addition,

3. $\forall z, z', z'' \in \mathcal{Z}, \rho(z', z'') \leq \rho(z, z') + \rho(z, z'')$,

$(\mathcal{Z}, \rho)$ is said to be a metric space and ρ is called a metric on $\mathcal{Z}$.

DEFINITION 6. (Negative type) The semimetric space $(\mathcal{Z}, \rho)$ is said to have negative type if $\forall n \geq 2, z_1, \dots, z_n \in \mathcal{Z}$, and $\alpha_1, \dots, \alpha_n \in \mathbb{R}$, with $\sum_{i=1}^n \alpha_i = 0$,

$$(2.7) \quad \sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j \rho(z_i, z_j) \leq 0.$$

Note that in the terminology of [Berg et al. \[1984\]](#), ρ satisfying (2.7) is said to be a *negative definite* function. The following proposition is derived from [\[Berg et al., 1984, Corollary 2.10, p. 78, and Proposition 3.2, p. 82\]](#).

PROPOSITION 7.

1. If ρ satisfies (2.7), then so does ρ^q , for $0 < q < 1$.
2. ρ is a semimetric of negative type if and only if there exists a Hilbert space $\mathcal{H}$ and an injective map $\varphi : \mathcal{Z} \rightarrow \mathcal{H}$, such that

$$(2.8) \quad \rho(z, z') = \|\varphi(z) - \varphi(z')\|_{\mathcal{H}}^2.$$

The second part of the Proposition shows that $(\mathbb{R}^d, \|\cdot - \cdot\|^2)$ is of negative type, and by taking $q = 1/2$ in the first part, we conclude that all Euclidean spaces are of negative type. In addition, whenever ρ is a semimetric of negative type, $\rho^{1/2}$ is a metric of negative type, i.e., even though ρ may not satisfy the triangle inequality, its square root must do if it obeys (2.7).

2.4. *Distance kernels.* Lyons [2011, p. 9] also uses the results in Proposition 7 to study embeddings to general Hilbert spaces, however the relation with the theory of reproducing kernel Hilbert spaces is not exploited. Semimetrics of negative type and symmetric positive definite kernels are in fact closely related, as summarized in the following Lemma, adapted from [Berg et al., 1984, Lemma 2.1, p. 74].

LEMMA 8. *Let $\mathcal{Z}$ be a nonempty set, and $\rho : \mathcal{Z} \times \mathcal{Z} \rightarrow \mathbb{R}$ a semimetric on $\mathcal{Z}$. Let $z_0 \in \mathcal{Z}$, and denote $k(z, z') = \rho(z, z_0) + \rho(z', z_0) - \rho(z, z')$. Then k is positive definite if and only if ρ satisfies (2.7).*

As a consequence, $k(z, z')$ defined above is a valid kernel on $\mathcal{Z}$ whenever ρ is a semimetric of negative type. For convenience, we will work with such kernels scaled by $1/2$.

DEFINITION 9. (**Distance-induced kernel**) Let ρ be a semimetric of negative type on $\mathcal{Z}$ and let $z_0 \in \mathcal{Z}$. The kernel

$$(2.9) \quad k(z, z') = \frac{1}{2} [\rho(z, z_0) + \rho(z', z_0) - \rho(z, z')]$$

is said to be the *distance-induced kernel* induced by ρ and *centred* at z_0 .

For brevity, we will drop “induced” hereafter, and say that k is simply the *distance kernel* (with some abuse of terminology). Note that distance kernels are not strictly positive definite, i.e., it is not true that $\forall n \in \mathbb{N}$, and for distinct $z_1, \dots, z_n \in \mathcal{Z}$,

$$\sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j k(z_i, z_j) = 0 \implies \alpha_i = 0 \forall i.$$

Indeed, if k were given by (2.9), it would suffice to take $n = 1$, since $k(z_0, z_0) = 0$. By varying the point at the center z_0 , we obtain a family

$$\mathcal{K}_\rho = \left\{ \frac{1}{2} [\rho(z, z_0) + \rho(z', z_0) - \rho(z, z')] \right\}_{z_0 \in \mathcal{Z}}$$

of distance kernels induced by ρ . We may now express (2.8) from Proposition

7 in terms of the canonical feature map for the RKHS $\mathcal{H}_k$.

PROPOSITION 10. *Let $(\mathcal{Z}, \rho)$ be a semimetric space of negative type, and $k \in \mathcal{K}_\rho$. Then:*

1. *k is nondegenerate, i.e., the Aronszajn map $z \mapsto k(\cdot, z)$ is injective.*
2. *$\rho(z, z') = k(z, z) + k(z', z') - 2k(z, z') = \|k(\cdot, z) - k(\cdot, z')\|_{\mathcal{H}_k}^2$.*

PROOF. If $z, z' \in \mathcal{Z}$ are such that $k(w, z) = k(w, z')$, for all $w \in \mathcal{Z}$, we would also have $\rho(z, z_0) - \rho(z, w) = \rho(z', z_0) - \rho(z', w)$, for all $w \in \mathcal{Z}$. In particular, by inserting $w = z$, and $w = z'$, we obtain $\rho(z, z') = -\rho(z, z') = 0$, i.e., $z = z'$. The second statement follows readily by expressing k in terms of ρ . $\square$

EXAMPLE 11. Let $\mathcal{Z} \subseteq \mathbb{R}^d$ and write $\rho_q(z, z') = \|z - z'\|^q$. By Proposition 7, ρ_q is a valid semimetric of negative type for $0 < q \leq 2$. The corresponding kernel centered at $z_0 = 0$ is given by:

$$(2.10) \quad k_q(z, z') = \frac{1}{2} (\|z\|^q + \|z'\|^q - \|z - z'\|^q).$$

2.5. *Semimetrics generated by kernels.* We now further develop the link between semimetrics of negative type and kernels. We start with a simple corollary of Proposition 7:

COROLLARY 12. *Let k be any nondegenerate kernel on $\mathcal{Z}$. Then,*

$$(2.11) \quad \rho(z, z') = k(z, z) + k(z', z') - 2k(z, z')$$

defines a valid semimetric ρ of negative type on $\mathcal{Z}$.

DEFINITION 13. **(Equivalent kernels)** Whenever the kernel k and semimetric ρ satisfy (2.11), we will say that k *generates* ρ . If two kernels generate the same semimetric, we will say that they are *equivalent kernels*.

It is clear that every distance kernel $\tilde{k} \in \mathcal{K}_\rho$ induced by ρ , also generates ρ . However, there are many other kernels that generate ρ , as illustrated in the following example.

EXAMPLE 14. Let $\mathcal{Z} = \mathbb{R}^d$, and consider the Gaussian kernel $k(z, z') = e^{-\sigma\|z-z'\|^2}$. The induced semimetric is $\rho(z, z') = 2 \left[1 - e^{-\sigma\|z-z'\|^2} \right]$. The Gaussian kernel is equivalent to $\tilde{k}(z, z') = e^{-\sigma\|z-z'\|^2} + 1 - e^{-\sigma\|z\|^2} - e^{-\sigma\|z'\|^2}$. Note that $\tilde{k} \in \mathcal{K}_\rho$, while $k \notin \mathcal{K}_\rho$.

PROPOSITION 15. Let k and $\tilde{k}$ be two kernels on $\mathcal{Z}$. k and $\tilde{k}$ are equivalent if and only if $\tilde{k}(z, z') = k(z, z') + f(z) + f(z')$, for some shift function $f : \mathcal{Z} \rightarrow \mathbb{R}$.

PROOF. Let $\tilde{k}(z, z') = k(z, z') + f(z) + f(z')$. Then,

$$\begin{aligned} \tilde{k}(z, z) + \tilde{k}(z', z') - 2\tilde{k}(z, z') &= k(z, z) + 2f(z) + k(z', z') + 2f(z') \\ &\quad - 2[k(z, z') + f(z) + f(z')] \\ &= k(z, z) + k(z', z') - 2k(z, z'). \end{aligned}$$

The converse is clear from writing $h = \tilde{k} - k$, whereby if k and $\tilde{k}$ are equivalent, it follows that $h(z, z') = \frac{1}{2}(h(z, z) + h(z', z'))$, for all $z, z' \in \mathcal{Z}$. $\square$

Not every choice of shift function f in Proposition 15 will be valid, as both k and $\tilde{k}$ are required to be positive definite. An important class of shift functions can be derived using RKHS functions, however. Namely, let k be a kernel on $\mathcal{Z}$ and let $f \in \mathcal{H}_k$, and define a kernel

$$\begin{aligned} \tilde{k}_f(z, z') &= \langle k(\cdot, z) - f, k(\cdot, z') - f \rangle_{\mathcal{H}_k} \\ &= k(z, z') - f(z) - f(z') + \|f\|_{\mathcal{H}_k}^2. \end{aligned}$$

Since it is representable as an inner product in a Hilbert space, $\tilde{k}_f$ is a valid kernel which is equivalent to k by Proposition 15. As a special case, if $f = \mu_k(P)$ for some $P \in \mathcal{M}_+^1(\mathcal{Z})$, we obtain the kernel centred at probability measure P :

$$(2.12) \quad \tilde{k}_P(z, z') := k(z, z') + \mathbb{E}_{WW'} k(W, W') - \mathbb{E}_W k(z, W) - \mathbb{E}_W k(z', W),$$

with $W, W' \stackrel{i.i.d.}{\sim} P$. Note that $\mathbb{E}_{ZZ' \stackrel{i.i.d.}{\sim} P} \tilde{k}_P(Z, Z') = 0$, i.e., $\mu_{\tilde{k}_P}(P) = 0$. The kernels of form (2.12) that are centred at the point masses $P = \delta_{z_0}$ are precisely the distance kernels equivalent to k .

The relationship between positive definite kernels and semimetrics of negative type is illustrated in Figure 1.




FIGURE 1. The relationship between kernels and semimetrics

REMARK 16. The requirement that kernels be characteristic (as introduced below Definition 3) is clearly important in hypothesis testing. A second family of kernels, widely used in the machine learning literature, are the universal kernels: universality can be used to guarantee consistency of learning algorithms [Steinwart and Christmann, 2008]. A relation between universal and characteristic kernels is described in Appendix B.

2.6. *Existence of kernel embedding through a semimetric.* In Section 2.1, we have seen that a sufficient condition for the kernel embedding $\mu_k(\nu)$ of $\nu \in \mathcal{M}(\mathcal{Z})$ to exist is that $\nu \in \mathcal{M}_k^{1/2}(\mathcal{Z})$. We will now interpret this condition in terms of the semimetric ρ generated by k .

DEFINITION 17. For $\theta > 0$, we say that $\nu \in \mathcal{M}(\mathcal{Z})$ has a finite θ -moment with respect to a semimetric ρ of negative type if there exists $z_0 \in \mathcal{Z}$, such that $\int \rho^\theta(z, z_0) d|\nu|(z) < \infty$. We denote

$$(2.13) \quad \mathcal{M}_\rho^\theta(\mathcal{Z}) = \{\nu \in \mathcal{M}(\mathcal{Z}) \quad : \quad \exists z_0 \in \mathcal{Z},$$

$$s.t. \quad \int \rho^\theta(z, z_0) d|\nu|(z) < \infty\}.$$

We now relate the space of measures with finite θ -moment with respect to ρ with space $\mathcal{M}_k^\theta(\mathcal{Z})$.

PROPOSITION 18. *Let k be a kernel that generates semimetric ρ , and let $n \in \mathbb{N}$. Then $\mathcal{M}_k^{n/2}(\mathcal{Z}) = \mathcal{M}_\rho^{n/2}(\mathcal{Z})$. In particular, if k_1 and k_2 generate the same semimetric ρ , then $\mathcal{M}_{k_1}^{n/2}(\mathcal{Z}) = \mathcal{M}_{k_2}^{n/2}(\mathcal{Z})$.*

PROOF. Let $\theta \geq \frac{1}{2}$. Suppose $\nu \in \mathcal{M}_k^\theta(\mathcal{Z})$. Then we have

$$\begin{aligned}
& \int \rho^\theta(z, z_0) d|\nu|(z) \\
&= \int \|k(\cdot, z) - k(\cdot, z_0)\|_{\mathcal{H}_k}^{2\theta} d|\nu|(z) \\
&\leq \int (\|k(\cdot, z)\|_{\mathcal{H}_k} + \|k(\cdot, z_0)\|_{\mathcal{H}_k})^{2\theta} d|\nu|(z) \\
&\leq 2^{2\theta-1} \left(\int \|k(\cdot, z)\|_{\mathcal{H}_k}^{2\theta} d|\nu|(z) + \int \|k(\cdot, z_0)\|_{\mathcal{H}_k}^{2\theta} d|\nu|(z) \right) \\
&= 2^{2\theta-1} \left(\int k^\theta(z, z) d|\nu|(z) + k^\theta(z_0, z_0) |\nu|(\mathcal{Z}) \right) \\
&< \infty,
\end{aligned}$$

where we have used that $a^{2\theta}$ is a convex function of a . From the above it is clear that $\mathcal{M}_k^\theta(\mathcal{Z}) \subset \mathcal{M}_\rho^\theta(\mathcal{Z})$ for $\theta \geq 1/2$.

To prove the other direction, we show by induction that $\mathcal{M}_\rho^\theta(\mathcal{Z}) \subset \mathcal{M}_k^{n/2}(\mathcal{Z})$ for $\theta \geq \frac{n}{2}$, $n \in \mathbb{N}$. Let $n = 1$. Let $\theta \geq \frac{1}{2}$, and suppose that $\nu \in \mathcal{M}_\rho^\theta(\mathcal{X})$. Then, by invoking the reverse triangle and Jensen's inequalities, we have:

$$\begin{aligned}
\int \rho^\theta(z, z_0) d|\nu|(z) &= \int \|k(\cdot, z) - k(\cdot, z_0)\|_{\mathcal{H}_k}^{2\theta} d|\nu|(z) \\
&\geq \int \left| k^{1/2}(z, z) - k^{1/2}(z_0, z_0) \right|^{2\theta} d|\nu|(z) \\
&\geq \left| \int k^{1/2}(z, z) d|\nu|(z) - \|\nu\|_{TV} k^{1/2}(z_0, z_0) \right|^{2\theta},
\end{aligned}$$

which implies $\nu \in \mathcal{M}_k^{1/2}(\mathcal{Z})$, thereby satisfying the result for $n = 1$. Suppose the result holds for $\theta \geq \frac{n-1}{2}$, i.e., $\mathcal{M}_\rho^\theta(\mathcal{Z}) \subset \mathcal{M}_k^{(n-1)/2}(\mathcal{Z})$ for $\theta \geq \frac{n-1}{2}$. Let $\nu \in \mathcal{M}_\rho^\theta(\mathcal{Z})$ for $\theta \geq \frac{n}{2}$. Then we have

$$\begin{aligned}
& \int \rho^\theta(z, z_0) d|\nu|(z) \\
&= \int (\|k(\cdot, z) - k(\cdot, z_0)\|_{\mathcal{H}_k}^n)^{\frac{2\theta}{n}} d|\nu|(z) \\
&\geq \left(\int \|k(\cdot, z) - k(\cdot, z_0)\|_{\mathcal{H}_k}^n d|\nu|(z) \right)^{\frac{2\theta}{n}} \\
&\geq \left(\int |\|k(\cdot, z)\|_{\mathcal{H}_k} - \|k(\cdot, z_0)\|_{\mathcal{H}_k}|^n d|\nu|(z) \right)^{\frac{2\theta}{n}} \\
&\geq \left| \int (\|k(\cdot, z)\|_{\mathcal{H}_k} - \|k(\cdot, z_0)\|_{\mathcal{H}_k})^n d|\nu|(z) \right|^{\frac{2\theta}{n}} \\
&= \left| \int \sum_{r=0}^n (-1)^r \binom{n}{r} \|k(\cdot, z)\|_{\mathcal{H}_k}^{n-r} \|k(\cdot, z_0)\|_{\mathcal{H}_k}^r d|\nu|(z) \right|^{\frac{2\theta}{n}} \\
&= \left| \underbrace{\int k^{\frac{n}{2}}(z, z) d|\nu|(z)}_A \right. \\
&\quad \left. + \underbrace{\sum_{r=1}^n (-1)^r \binom{n}{r} k^{\frac{r}{2}}(z_0, z_0) \int k^{\frac{n-r}{2}}(z, z) d|\nu|(z)}_B \right|^{\frac{2\theta}{n}}.
\end{aligned}$$

Note that the terms in B are finite since for $\theta \geq \frac{n}{2} \geq \frac{n-1}{2} \geq \dots \geq \frac{1}{2}$, we have $\mathcal{M}_\rho^\theta(\mathcal{Z}) \subset \mathcal{M}_k^{(n-1)/2}(\mathcal{Z}) \subset \dots \subset \mathcal{M}_k^1(\mathcal{Z}) \subset \mathcal{M}_k^{1/2}(\mathcal{Z})$ and therefore A is finite, which means $\nu \in \mathcal{M}_k^{n/2}(\mathcal{Z})$, i.e., $\mathcal{M}_\rho^\theta(\mathcal{Z}) \subset \mathcal{M}_k^{n/2}(\mathcal{Z})$ for $\theta \geq \frac{n}{2}$. The result shows that $\mathcal{M}_\rho^\theta(\mathcal{Z}) = \mathcal{M}_k^\theta(\mathcal{Z})$ for all $\theta \in \{\frac{n}{2} : n \in \mathbb{N}\}$. $\square$

The above Proposition gives a natural interpretation of conditions on probability measures in terms of moments w.r.t. ρ . Namely, the kernel embedding $\mu_k(P)$, where kernel k generates the semimetric ρ , exists for every P with finite half-moment w.r.t. ρ , and thus the MMD $\gamma_k(P, Q)$ between P and Q is well defined whenever both P and Q have finite half-moments w.r.t. ρ . Furthermore, HSIC between random variables X and Y is well defined whenever their marginals P_X and P_Y have finite first moments w.r.t. semimetric ρ_X and ρ_Y generated by kernels k_X and k_Y on their respective domains $\mathcal{X}$ and $\mathcal{Y}$.

3. Equivalence of MMD and energy distance. In this section, we begin with a review of the energy distance, which measures distance between distributions, and demonstrate that it is an instance of the maximum mean discrepancy introduced in Definition 3. The energy distance was intro-

duced by Székely and Rizzo [2004, 2005] as a measure of statistical distance between two probability measures P and Q on $\mathbb{R}^d$ with finite first moments,

$$(3.1) \quad D_E(P, Q) = 2\mathbb{E}_{ZW} \|Z - W\| - \mathbb{E}_{ZZ'} \|Z - Z'\| - \mathbb{E}_{WW'} \|W - W'\|,$$

where $Z, Z' \stackrel{i.i.d.}{\sim} P$ and $W, W' \stackrel{i.i.d.}{\sim} Q$. This quantity is always nonnegative, and is strictly positive if $P \neq Q$. In scalar case, it coincides with twice the Cramer-Von Mises distance.

Following Lyons [2011], the notion is readily generalized to a semimetric space of negative type.

DEFINITION 19. Let $(\mathcal{Z}, \rho)$ be a semimetric space of negative type, and let $P, Q \in \mathcal{M}_\rho^1(\mathcal{Z})$. The energy distance between P and Q , w.r.t. ρ is

$$(3.2) \quad D_E(P, Q) = 2\mathbb{E}_{ZW} \rho(Z, W) - \mathbb{E}_{ZZ'} \rho(Z, Z') - \mathbb{E}_{WW'} \rho(W, W'),$$

where $Z, Z' \stackrel{i.i.d.}{\sim} P$ and $W, W' \stackrel{i.i.d.}{\sim} Q$.

The moment condition is required in order to ensure that each of the expectations in (3.2) is finite. Note that the energy distance can equivalently be represented in the integral form,

$$(3.3) \quad D_{E,\rho}(P, Q) = - \int \int \rho d([P - Q] \times [P - Q]),$$

whereby the negative type of ρ implies the non-negativity of $D_{E,\rho}$.

We are now able to show that for every ρ , $D_{E,\rho}$ is related to the MMD associated to a kernel k that generates ρ .

THEOREM 20. Let $(\mathcal{Z}, \rho)$ be a semimetric space of negative type and let k be any kernel that generates ρ . Then

$$D_{E,\rho}(P, Q) = 2\gamma_k^2(P, Q), \quad \forall P, Q \in \mathcal{M}_+^1(\mathcal{Z}) \cap \mathcal{M}_\rho^1(\mathcal{Z}).$$

In particular, equivalent kernels have the same maximum mean discrepancy.

PROOF. Since k generates ρ , we can write $\rho(z, w) = k(z, z) + k(w, w) - 2k(z, w)$. Denote $\nu = P - Q$. Then,

$$\begin{aligned} D_{E,\rho}(P, Q) &= - \int \int \rho(z, w) d\nu(z) d\nu(w) \\ &= - \int \int [k(z, z) + k(w, w) - 2k(z, w)] d\nu(z) d\nu(w) \\ &= 2 \int \int k(z, w) d\nu(z) d\nu(w) \\ &= 2\gamma_k^2(P, Q). \end{aligned}$$

where we used the fact that $\nu(\mathcal{Z}) = 0$. $\square$

In its basic form, this result also appears in [Lyons, 2011, p. 11, Eq. (3.9)], but since Lyons works with embeddings into general Hilbert spaces, the link to the RKHS-based statistics and MMD in particular is obscured. Theorem 20 shows that all kernels that generate the same semimetric ρ on $\mathcal{Z}$ give rise to the same metric γ_k on (possibly a subset of) $\mathcal{M}_+^1(\mathcal{Z})$, whence γ_k is merely an extension of metric induced by $\rho^{1/2}$ on the point masses, since

$$\begin{aligned}\gamma_k(\delta_z, \delta_{z'}) &= \|k(\cdot, z) - k(\cdot, z')\|_{\mathcal{H}_k} \\ &= \rho^{1/2}(z, z').\end{aligned}$$

In other words, whenever kernel k generates ρ , $z \mapsto \delta_z$ is an isometry between $(\mathcal{Z}, \rho^{1/2})$ and $\{\delta_z : z \in \mathcal{Z}\} \subset \mathcal{M}_+^1(\mathcal{Z})$, endowed with the MMD metric $\gamma_k = \frac{1}{2}D_{E, \rho}^{1/2}$; and the Aronszajn map $z \mapsto k(\cdot, z)$ is an isometric embedding of a metric space $(\mathcal{Z}, \rho^{1/2})$ into $\mathcal{H}_k$. These isometries are depicted in Figure 2. For simplicity, we show the case of a bounded kernel, where kernel embeddings are well defined for all $P \in \mathcal{M}_+^1(\mathcal{Z})$, in which case $(\mathcal{M}_+^1(\mathcal{Z}), \gamma_k)$ and $\mu_k(\mathcal{M}_+^1(\mathcal{Z})) = \{\mu_k(P) : P \in \mathcal{M}_+^1(\mathcal{Z})\}$ endowed with the Hilbert-space metric inherited from $\mathcal{H}_k$ are also isometric (note that this implies that the corresponding subsets of RKHSs corresponding to equivalent kernels are also isometric).

REMARK 21. Theorem 20 requires that $P, Q \in \mathcal{M}_\rho^1(\mathcal{Z})$, i.e., that P and Q have finite first moments w.r.t. ρ , as otherwise the energy distance between P and Q may be undefined; e.g., each of the expectations $\mathbb{E}_{ZZ'}\rho(Z, Z')$, $\mathbb{E}_{WW'}\rho(W, W')$ and $\mathbb{E}_{ZW}\rho(Z, W)$ may be infinite. However, as long as a *weaker condition* $P, Q \in \mathcal{M}_\rho^{1/2}(\mathcal{Z})$ is satisfied, i.e., P and Q have finite *half*-moments w.r.t. ρ , the maximum mean discrepancy γ_k will be well defined. If, in addition, $P, Q \in \mathcal{M}_\rho^1(\mathcal{Z})$, then the energy distance between P and Q is also well defined, and must be equal to γ_k . We will later invoke the same condition $P, Q \in \mathcal{M}_k^1(\mathcal{Z})$ when describing the asymptotic distribution of the empirical maximum mean discrepancy in Section 6.

4. Distance covariance through kernel embeddings. A related notion to the energy distance is that of distance covariance, which measures dependence between random variables. In this section, we review distance covariance and show that it is an instance of the Hilbert-Schmidt independence criterion.




FIGURE 2. Isometries relating the semimetric ρ on $\mathcal{Z}$ with the RKHS corresponding to a kernel k that generates ρ , and with the set of probability measures on $\mathcal{Z}$.

4.1. *Distance covariance.* Let X be a random vector on $\mathbb{R}^p$ and Y a random vector on $\mathbb{R}^q$. The distance covariance was introduced by Székely et al. [2007], Székely and Rizzo [2009] to address the problem of testing and measuring dependence between X and Y in terms of a weighted L_2 -distance between characteristic functions of the joint distribution of X and Y and the product of their marginals. As a particular choice of weight function is used (we discuss this further in Section 4.3), it can be computed in terms of certain expectations of pairwise Euclidean distances,

$$(4.1) \quad \mathcal{V}^2(X, Y) = \mathbb{E}_{XY} \mathbb{E}_{X'Y'} \|X - X'\| \|Y - Y'\| \\ + \mathbb{E}_X \mathbb{E}_{X'} \|X - X'\| \mathbb{E}_Y \mathbb{E}_{Y'} \|Y - Y'\| \\ - 2 \mathbb{E}_{XY} [\mathbb{E}_{X'} \|X - X'\| \mathbb{E}_{Y'} \|Y - Y'\|],$$

where (X, Y) and (X', Y') are $\stackrel{i.i.d.}{\sim} P_{XY}$. As in the case of the energy distance, Lyons [2011] established that the generalization of the distance covariance is possible to metric space of negative type.

DEFINITION 22. Let $(\mathcal{X}, \rho_X)$ and $(\mathcal{Y}, \rho_Y)$ be semimetric spaces of negative type, and let $X \sim P_X \in \mathcal{M}_{\rho_X}^2(\mathcal{X})$ and $Y \sim P_Y \in \mathcal{M}_{\rho_Y}^2(\mathcal{Y})$, having joint

distribution P_{XY} . The generalized distance covariance of X and Y is

$$(4.2) \quad \mathcal{V}^2(X, Y) = \mathbb{E}_{XY} \mathbb{E}_{X'Y'} \rho_X(X, X') \rho_Y(Y, Y') \\ + \mathbb{E}_X \mathbb{E}_{X'} \rho_X(X, X') \rho_Y(Y, Y') \\ - 2 \mathbb{E}_{XY} [\mathbb{E}_{X'} \rho_X(X, X') \mathbb{E}_{Y'} \rho_Y(Y, Y')].$$

As with the MMD, the moment condition ensures that the expectations are finite (which can be seen using Cauchy-Schwarz inequality). Equivalently, the generalized distance covariance can be represented in integral form,

$$(4.3) \quad \mathcal{V}_{\rho_X, \rho_Y}^2(X, Y) = \int \int \rho_X \rho_Y d([P_{XY} - P_X P_Y] \times [P_{XY} - P_X P_Y]),$$

where $\rho_X \rho_Y$ is viewed as a function on $(\mathcal{X} \times \mathcal{Y}) \times (\mathcal{X} \times \mathcal{Y})$. Furthermore, Lyons [2011, Theorem 3.20] shows that distance covariance in a metric space characterizes independence (i.e., $\mathcal{V}_{\rho_X, \rho_Y}^2(X, Y) = 0$ if and only if X and Y are independent) if the metrics ρ_X and ρ_Y satisfy an additional property, termed *strong negative type*. The discussion of this property is relegated to Section 5.

4.2. *Equivalence between HSIC and distance covariance.* The following theorem demonstrates the link between HSIC and the distance covariance.

THEOREM 23. *Let $(\mathcal{X}, \rho_X)$ and $(\mathcal{Y}, \rho_Y)$ be semimetric spaces of negative type, and let $X \sim P_X \in \mathcal{M}_{\rho_X}^2(\mathcal{X})$ and $Y \sim P_Y \in \mathcal{M}_{\rho_Y}^2(\mathcal{Y})$, having joint distribution P_{XY} . Let k_X and k_Y be any two kernels on $\mathcal{X}$ and $\mathcal{Y}$ that generate ρ_X and ρ_Y , respectively, and denote*

$$(4.4) \quad k((x, y), (x', y')) = k_X(x, x') k_Y(y, y').$$

Then, $\mathcal{V}_{\rho_X, \rho_Y}^2(X, Y) = 4\gamma_k^2(P_{XY}, P_X P_Y)$.

PROOF. Put $\nu = P_{XY} - P_X P_Y$. Then

$$\begin{aligned} \mathcal{V}_{\rho_X, \rho_Y}^2(X, Y) &= \int \int \rho_X(x, x') \rho_Y(y, y') d\nu(x, y) d\nu(x', y') \\ &= \int \int (k_X(x, x) + k_X(x', x') - 2k_X(x, x')) \cdot \\ &\quad (k_Y(y, y) + k_Y(y', y') - 2k_Y(y, y')) d\nu(x, y) d\nu(x', y') \\ &= 4 \int \int k_X(x, x') k_Y(y, y') d\nu(x, y) d\nu(x', y') \\ &= 4\gamma_k^2(P_{XY}, P_X P_Y), \end{aligned}$$

where we used that $\nu(\mathcal{X} \times \mathcal{Y}) = 0$, and that $\int g(x, y, x', y') d\nu(x, y) d\nu(x', y') = 0$ when g does not depend on one or more of its arguments, since ν also has zero marginal measures. Convergence of integrals of the form $\int k_{\mathcal{X}}(x, x) k_{\mathcal{Y}}(y, y) d\nu(x, y)$ is ensured by the moment conditions on the marginals. $\square$

We remark that a similar result to Theorem 23 is given by Lyons [2011, Proposition 3.16], but without making use of the link with kernel embeddings. Theorem 23 is a more general statement, in the sense that we allow ρ to be a semimetric of negative type, rather than metric. In addition, the kernel interpretation leads to a significantly simpler proof: the result is an immediate application of the HSIC expansion in (2.6).

REMARK 24. As in Remark 21, to ensure the existence of the distance covariance, we impose a stronger condition on the marginals: $P_X \in \mathcal{M}_{k_X}^2(\mathcal{X})$ and $P_Y \in \mathcal{M}_{k_Y}^2(\mathcal{Y})$, while $P_X \in \mathcal{M}_{k_X}^1(\mathcal{X})$ and $P_Y \in \mathcal{M}_{k_Y}^1(\mathcal{Y})$ are sufficient for the existence of the Hilbert-Schmidt independence criterion.

REMARK 25. As introduced by Székely et al. [2007], the notion of distance covariance extends naturally to that of *distance variance* $\mathcal{V}^2(X) = \mathcal{V}^2(X, X)$ and of *distance correlation* (by analogy with the Pearson product-moment correlation coefficient),

$$\mathcal{R}^2(X, Y) = \begin{cases} \frac{\mathcal{V}^2(X, Y)}{\mathcal{V}(X)\mathcal{V}(Y)}, & \mathcal{V}(X)\mathcal{V}(Y) > 0, \\ 0, & \mathcal{V}(X)\mathcal{V}(Y) = 0. \end{cases}$$

The distance correlation can also be expressed in terms of associated kernels - see Appendix A for details.

4.3. *Characteristic function interpretation.* The distance covariance in (4.1) was defined by [Székely et al., 2007] in terms of a weighted distance between characteristic functions. We briefly review this interpretation here, however we show that this approach *cannot* be used to derive a kernel-based measure of dependence (this result was first noted by Gretton et al. [2009b], and is included here in the interest of completeness). Let X be a random vector on $\mathcal{X} = \mathbb{R}^p$ and Y a random vector on $\mathcal{Y} = \mathbb{R}^q$. The characteristic function of X and Y , respectively, will be denoted by f_X and f_Y , and their joint characteristic function by f_{XY} . The distance covariance $\mathcal{V}(X, Y)$ is defined via the norm of $f_{XY} - f_X f_Y$ in a weighted L_2 space on $\mathbb{R}^{p+q}$, i.e.,

$$(4.5) \quad \mathcal{V}^2(X, Y) = \int_{\mathbb{R}^{p+q}} |f_{X,Y}(t, s) - f_X(t) f_Y(s)|^2 w(t, s) dt ds,$$

for a particular choice of weight function given by

$$(4.6) \quad w(t, s) = \frac{1}{c_p c_q} \cdot \frac{1}{\|t\|^{1+p} \|s\|^{1+q}},$$

where $c_d = \pi^{\frac{1+d}{2}} / \Gamma(\frac{1+d}{2})$, $d \geq 1$. An important aspect of distance covariance is that $\mathcal{V}(X, Y) = 0$ if and only if X and Y are independent. We next obtain a similar statistic in the kernel setting. Write $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$, and let $k(z, z') = \kappa(z - z')$ be a translation invariant RKHS kernel on $\mathcal{Z}$, where $\kappa : \mathcal{Z} \rightarrow \mathbb{R}$ is a bounded continuous function. Using Bochner's theorem, κ can be written as:

$$\kappa(z) = \int e^{-z^\top u} d\Lambda(u),$$

for a finite non-negative Borel measure Λ . It follows [Gretton et al., 2009b] that

$$\gamma_k^2(P_{XY}, P_X P_Y) = \int_{\mathbb{R}^{p+q}} |f_{X,Y}(t, s) - f_X(t) f_Y(s)|^2 d\Lambda(t, s),$$

which is in clear correspondence with (4.5). However, the weight function in (4.6) is not integrable — so we cannot find a continuous translation invariant kernel for which γ_k coincides with the distance covariance. By contrast, note the kernel in (4.4) is *not* translation invariant.

5. Distinguishing probability distributions. Theorem 3.20 of Lyons [2011] shows that distance covariance in a metric space characterizes independence if the metrics satisfy an additional property, termed *strong negative type*. We will extend this notion to a semimetric ρ , establish the interpretation of strong negative type in terms of kernels, and show how the strong negative type of a semimetric ρ can be established by considering whether the kernel k that generates ρ is characteristic.

DEFINITION 26. The semimetric space $(\mathcal{Z}, \rho)$, where ρ is generated by kernel k , is said to have a *strong negative type* if $\forall P, Q \in \mathcal{M}_+^1(\mathcal{Z}) \cap \mathcal{M}_k^1(\mathcal{Z})$,

$$(5.1) \quad P \neq Q \Rightarrow \int \rho d([P - Q] \times [P - Q]) < 0.$$

Since the quantity in (5.1) is exactly $-2\gamma_k^2(P, Q)$, $\forall P, Q \in \mathcal{M}_+^1(\mathcal{Z}) \cap \mathcal{M}_k^1(\mathcal{Z})$, we directly obtain:

PROPOSITION 27. *Let kernel k generate ρ . Then $(\mathcal{Z}, \rho)$ has a strong negative type if and only if k is characteristic to $\mathcal{M}_+^1(\mathcal{Z}) \cap \mathcal{M}_k^1(\mathcal{Z})$.*

Thus, the problems of checking whether a semimetric is of strong negative type and whether its associated kernel is characteristic to an appropriate space of Borel probability measures are equivalent. This conclusion has some overlap with [Lyons, 2011]: in particular, Proposition 27 is stated in [Lyons, 2011, Proposition 3.10], where the barycenter map β is a kernel embedding in our terminology, although Lyons does not consider distribution embeddings in an RKHS.

REMARK 28. From [Lyons, 2011, Theorem 3.25], every separable Hilbert space $\mathcal{Z}$ is of strong negative type, so a distance kernel k induced by the (inner product) metric on $\mathcal{Z}$ is characteristic to the appropriate space of probability measures.

REMARK 29. Consider the kernel in (4.4), and let us, for simplicity, assume that $k_{\mathcal{X}}$ and $k_{\mathcal{Y}}$ are bounded, so that we can consider embeddings of all probability measures. It turns out that k need not be characteristic - i.e., it may not be able to distinguish between any two distributions on $\mathcal{X} \times \mathcal{Y}$, even if $k_{\mathcal{X}}$ and $k_{\mathcal{Y}}$ are characteristic. Namely, if $k_{\mathcal{X}}$ is the distance kernel induced by $\rho_{\mathcal{X}}$ and centred at x_0 , then $k((x_0, y), (x_0, y')) = 0$ for all $y, y' \in \mathcal{Y}$. That means that for every two distinct $P_Y, Q_Y \in \mathcal{M}_+^1(\mathcal{Y})$, we have $\gamma_k^2(\delta_{x_0} P_Y, \delta_{x_0} Q_Y) = 0$. Thus, given that $\rho_{\mathcal{X}}$ and $\rho_{\mathcal{Y}}$ have strong negative type, the kernel in (4.4) characterizes independence, but not equality of probability measures on the product space. Informally speaking, distinguishing P_{XY} from $P_X P_Y$ is an easier problem than two-sample testing on the product space.

6. Empirical estimates and hypothesis tests.

6.1. *Two-sample testing.* So far, we have seen that the population expression of the MMD between P and Q is well defined as long as P and Q lie in the space $\mathcal{M}_k^{1/2}(\mathcal{Z})$, or, equivalently, have a finite half-moment w.r.t. semimetric ρ generated by k . However, this assumption will not suffice to establish a meaningful hypothesis test using empirical estimates of the MMD. We will require a stronger condition, that $P, Q \in \mathcal{M}_+^1(\mathcal{Z}) \cap \mathcal{M}_k^1(\mathcal{Z})$ (which is the same condition under which the energy distance is well defined). Note that, under this condition we also have $k \in L_{P \times P}^2(\mathcal{Z} \times \mathcal{Z})$, as $\int \int k^2(z, z') dP(z) dP(z') \leq \left(\int k(z, z) dP(z) \right)^2$.

Given i.i.d. samples $\mathbf{z} = \{z_i\}_{i=1}^m \sim P$ and $\mathbf{w} = \{w_i\}_{i=1}^n \sim Q$, the empirical (biased) V-statistic estimate of (2.3) is given by:

$$\begin{aligned}
\hat{\gamma}_{k,V}^2(\mathbf{Z}, \mathbf{W}) &= \gamma_k^2 \left(\frac{1}{m} \sum_{i=1}^m \delta_{z_i}, \frac{1}{n} \sum_{j=1}^n \delta_{w_j} \right) \\
&= \frac{1}{m^2} \sum_{i=1}^m \sum_{j=1}^m k(z_i, z_j) + \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n k(w_i, w_j) \\
&\quad - \frac{2}{mn} \sum_{i=1}^m \sum_{j=1}^n k(z_i, w_j).
\end{aligned}
\tag{6.1}$$

Recall that if k generates ρ , this estimate involves only the pairwise ρ -distances between the sample points.

We now describe a two-sample test using this statistic. The kernel $\tilde{k}_P$ centred at P in (2.12) plays a key role in characterizing the null distribution of degenerate V-statistic. To $\tilde{k}_P$, we associate the *integral kernel operator* $S_{\tilde{k}_P} : L_P^2(\mathcal{Z}) \rightarrow L_P^2(\mathcal{Z})$ (cf. e.g., [Steinwart and Christmann, 2008, p.126-127]), given by:

$$S_{\tilde{k}_P} g(z) = \int_{\mathcal{Z}} \tilde{k}_P(z, w) g(w) dP(w).$$

The condition that $P \in \mathcal{M}_k^1(\mathcal{Z})$, and, as a consequence, that $\tilde{k}_P \in L_{P \times P}^2(\mathcal{Z} \times \mathcal{Z})$, is closely related to the desired properties of the integral operator. Namely, this implies that $S_{\tilde{k}_P}$ is a trace class operator, and, thus, a Hilbert-Schmidt operator [Reed and Simon, 1980, Proposition VI.23]. The following theorem is a special case of [Gretton et al., 2012, Theorem 12], where for simplicity, we focus on the case where $m = n$.

THEOREM 30. *Let k be a kernel on $\mathcal{Z}$, and let $\mathbf{Z} = \{Z_i\}_{i=1}^m$ and $\mathbf{W} = \{W_i\}_{i=1}^m$ be two i.i.d. samples from $P \in \mathcal{M}_+^1(\mathcal{Z}) \cap \mathcal{M}_k^1(\mathcal{Z})$. Then*

$$\frac{m}{2} \hat{\gamma}_{k,V}^2(\mathbf{Z}, \mathbf{W}) \rightsquigarrow \sum_{i=1}^{\infty} \lambda_i N_i^2,$$

where $N_i \stackrel{i.i.d.}{\sim} \mathcal{N}(0, 1)$, $i \in \mathbb{N}$, and $\{\lambda_i\}_{i=1}^{\infty}$ are the eigenvalues of the operator $S_{\tilde{k}_P}$.

Note that the limiting expression in (6.3) is a valid random variable precisely since $S_{\tilde{k}_P}$ is Hilbert-Schmidt, i.e., since $\sum_{i=1}^{\infty} \lambda_i^2 < \infty$.

6.2. *Independence testing.* In the case of independence testing, we are given i.i.d. samples $\mathbf{z} = \{(x_i, y_i)\}_{i=1}^m \sim P_{XY}$, and the resulting V-statistic estimate (HSIC) is [Gretton et al., 2005, 2008]:

$$(6.4) \quad \text{HSIC}(\mathbf{z}; k_X, k_Y) = \frac{1}{m^2} \text{Tr}(K_X H K_Y H),$$

where K_X , K_Y and H are $m \times m$ matrices given by $(K_X)_{ij} := k_X(x_i, x_j)$, $(K_Y)_{ij} := k_Y(y_i, y_j)$ and $H_{ij} = \delta_{ij} - \frac{1}{m}$ (centering matrix). The null distribution of HSIC takes an analogous form to (6.3) of a weighted sum of chi-squares, but with coefficients corresponding to the products of the eigenvalues of integral operators $S_{\tilde{k}_{P_X}} : L_{P_X}^2(\mathcal{X}) \rightarrow L_{P_X}^2(\mathcal{X})$, and $S_{\tilde{k}_{P_Y}} : L_{P_Y}^2(\mathcal{Y}) \rightarrow L_{P_Y}^2(\mathcal{Y})$. Similarly to the case of two-sample testing, we will require that $P_X \in \mathcal{M}_{k_X}^1(\mathcal{X})$ and $P_Y \in \mathcal{M}_{k_Y}^1(\mathcal{Y})$, implying that integral operators $S_{\tilde{k}_{P_X}}$ and $S_{\tilde{k}_{P_Y}}$ are trace class operators. The following Theorem is from [Zhang et al., 2011, Theorem 4]. See also [Lyons, 2011, Remark 2.9].

THEOREM 31. *Let $\mathbf{Z} = \{(X_i, Y_i)\}_{i=1}^m$ be an i.i.d. sample from $P_{XY} = P_X P_Y$, with values in $\mathcal{X} \times \mathcal{Y}$, s.t. $P_X \in \mathcal{M}_{k_X}^1(\mathcal{X})$ and $P_Y \in \mathcal{M}_{k_Y}^1(\mathcal{Y})$. Then*

$$(6.5) \quad m\text{HSIC}(\mathbf{Z}; k_X, k_Y) \rightsquigarrow \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \lambda_i \eta_j N_{i,j}^2,$$

where $N_{i,j} \sim \mathcal{N}(0, 1)$, $i, j \in \mathbb{N}$, are independent and $\{\lambda_i\}_{i=1}^{\infty}$ and $\{\eta_j\}_{j=1}^{\infty}$ are the eigenvalues of the operators $S_{\tilde{k}_{P_X}}$ and $S_{\tilde{k}_{P_Y}}$, respectively.

6.3. *Test designs.* We would like to design distance-based tests with an asymptotic Type I error of α , and thus we require an estimate of the $(1 - \alpha)$ -quantile of the null distribution. We investigate two approaches, both of which yield consistent tests: a bootstrap approach [Arcones and Giné, 1992], and a spectral approach [Gretton et al., 2009a, Zhang et al., 2011]. The latter requires empirical computation of eigenvalues of kernel integral operators, a problem studied extensively in the context of kernel PCA [Schölkopf et al., 1997]. To estimate limiting distribution in (6.3), we compute the spectrum of the centred Gram matrix $\tilde{K} = H K H$ on the aggregated samples. Here, K is a $2m \times 2m$ matrix, with entries $K_{ij} = k(u_i, u_j)$, $\mathbf{u} = [\mathbf{z} \ \mathbf{w}]$ is the concatenation of the two samples and H is the centering matrix. Gretton et al. [2009a] show that the null distribution defined using the finite sample estimates of these eigenvalues converges to the population distribution, provided that the spectrum is square-root summable. As demonstrated in [Gretton et al.,

2009a], spectral estimation of the test threshold has a smaller computational cost of $\mathcal{O}(m^3)$ than that of the bootstrap-based approach, which is $\mathcal{O}(m^4)$, while providing an indistinguishable performance. The same approach can be used in obtaining a consistent finite sample null distribution for HSIC, via computation of the empirical eigenvalues of $\tilde{K}_X = HK_XH$ and $\tilde{K}_Y = HK_YH$; see [Zhang et al., 2011].

Both Székely and Rizzo [2004, p. 14] and Székely et al. [2007, p. 2782-2783] establish that the energy distance and distance covariance statistics, respectively, converge to the weighted sums of chi-squares of forms similar to (6.3). Analogous results for the generalized distance covariance are presented in [Lyons, 2011, p. 7-8]. These works do not propose test designs that attempt to estimate the coefficients λ_i , $i \in \mathbb{N}$, however. Besides the bootstrap, Székely et al. [2007, Theorem 6] also propose an independence test using a bound applicable to a general quadratic form Q of centered Gaussian random variables with $\mathbb{E}[Q] = 1$: $\mathbb{P}\{Q \geq (\Phi^{-1}(1 - \alpha/2))^2\} \leq \alpha$, valid for $0 < \alpha \leq 0.215$. When applied to the distance covariance statistic, the upper bound of α is achieved if X and Y are independent Bernoulli variables. The authors remark that the resulting criterion might be over-conservative. Thus, more sensitive distance covariance tests are possible by computing the spectrum of the centred Gram matrices associated to distance kernels, and we pursue this approach in the next section.

7. Experiments.

7.1. Two-sample experiments. In the two-sample experiments, we investigate three different kinds of synthetic data. In the first, we compare two multivariate Gaussians, where the means differ in one dimension only, and all variances are equal. In the second, we again compare two multivariate Gaussians, but this time with identical means in all dimensions, and variance that differs in a single dimension. In our third experiment, we use the benchmark data of Sriperumbudur et al. [2009]: one distribution is a univariate Gaussian, and the second is a univariate Gaussian with a sinusoidal perturbation of increasing frequency (where higher frequencies correspond to harder problems). All tests use a distance kernel induced by the Euclidean distance. As shown on the left hand plots in Figure 3, the spectral and bootstrap test designs appear indistinguishable, and significantly outperform the test designed using the quadratic form bound, which appears to be far too conservative for the data sets considered. This is confirmed by checking the Type I error of the quadratic form test, which is significantly smaller than the desired test size of $\alpha = 0.05$.




FIGURE 3. (left) MMD using Gaussian and distance kernels for various tests; (right) Spectral MMD using distance kernels with various exponents.

We also compare the performance to that of the Gaussian kernel, with the bandwidth set to the median distance between points in the aggregation of samples. We see that when the means differ, both tests perform similarly.

When the variances differ, it is clear that the Gaussian kernel has a major advantage over the distance kernel, although this advantage decreases with increasing dimension (where both perform poorly). In the case of a sinusoidal perturbation, the performance is again very similar.

In addition, following Example 11, we investigate performance of kernels obtained using the semimetric $\rho(z, z') = \|z - z'\|^q$ for $0 < q \leq 2$. Results are presented in the right hand plots of Figure 3. While we note that the judiciously chosen values of q offer some improvement in the cases of differing mean and variance, a dramatic improvement compared to the case $q = 1$ and the Gaussian kernel is noticeable in the case of sinusoidal perturbation, where the values $q = 1/3$ (and smaller) offer virtually error-free performance even at high frequencies (note that $q = 1$ yields the energy distance described in [Székely and Rizzo, 2004, 2005]).

We observe from the simulation results that distance kernels with higher exponents are advantageous in cases where distributions differ in mean value along a single dimension (with noise in the remainder), whereas distance kernels with smaller exponents are more sensitive to differences in distributions at finer lengthscales (i.e., where the characteristic functions of the distributions differ at higher frequencies).

7.2. Independence experiments. To assess independence tests, we used an artificial benchmark proposed by Gretton et al. [2008]: we generated univariate random variables from the ICA benchmark densities of Bach and Jordan [2002]; rotated them in the product space by an angle between 0 and $\pi/4$ to introduce dependence; filled additional dimensions with independent Gaussian noise; and, finally, passed the resulting multivariate data through random and independent orthogonal transformations. The resulting random variables X and Y were dependent but uncorrelated. The case $m = 128$ (sample size) and $d = 2$ (dimension) is plotted in Figure 4 (left). As observed by Gretton et al. [2009b], the Gaussian kernel does better than the distance kernel with $q = 1$. By varying q , however, we are able to obtain a wide performance range: in particular, the values $q = 1/3$ (and smaller) have an advantage over the Gaussian kernel on this dataset. As for the two-sample case, bootstrap and spectral tests have indistinguishable performance, and are significantly more sensitive than the quadratic form-based test, which failed to detect any dependence on this dataset.

In addition, we assess the performance on sinusoidally dependent data. The sample of the random variable pair X, Y was drawn from $P_{XY} \propto 1 + \sin(\ell x) \sin(\ell y)$ for integer ℓ , on the support $\mathcal{X} \times \mathcal{Y}$, where $\mathcal{X} := [-\pi, \pi]$ and $\mathcal{Y} := [-\pi, \pi]$. In this way, increasing ℓ causes the departure from a uniform

(independent) distribution to occur at increasing frequencies, making this departure harder to detect given a small sample size. Results are in Figure 4 (right). The distance covariance outperforms the Gaussian kernel on this example, and smaller exponents result in better performance (lower Type II error when the departure from independence occurs at higher frequencies). Finally, we note that the setting $q = 1$, as described by Székely et al. [2007], Székely and Rizzo [2009], is a reasonable heuristic in practice, but does not yield the most powerful tests on either dataset.




FIGURE 4. *HSIC using distance kernels with various exponents and a Gaussian kernel as a function of (left) the angle of rotation for the dependence induced by rotation; (right) frequency ℓ in the sinusoidal dependence example.*

8. Conclusion. We have established an equivalence between the generalized notions of energy distance and distance covariance, computed with respect to semimetrics of negative type, and RKHS measures of distance between distributions. In particular, energy distances and RKHS distance measures coincide when the semimetrics and kernels are related in a particular way.

The interpretation of the energy distance and distance covariance in an RKHS setting should be of considerable interest both to statisticians and machine learning researchers, since the associated kernels may be used much more widely: in conditional dependence testing and estimates of the chi-squared distance [Fukumizu et al., 2008], in Bayesian inference [Fukumizu et al., 2011], in mixture density estimation [Sriperumbudur, 2011] and in other machine learning applications. In particular, the link with kernels makes these applications of the energy distance immediate and straightforward. Finally, for problem settings defined most naturally in terms of distances, and where

these distances are of negative type, there is an interpretation in terms of reproducing kernels, and the learning machinery from the kernel literature can be brought to bear.

References.

- M. Arcones and E. Giné. On the bootstrap of U and V statistics. *Ann. Stat.*, 20(2): 655–674, 1992.
- F. R. Bach and M. I. Jordan. Kernel independent component analysis. *J. Mach. Learn. Res.*, 3:1–48, 2002.
- C. Berg, J. P. R. Christensen, and P. Ressel. *Harmonic Analysis on Semigroups*. Springer, New York, 1984.
- A. Berline and C. Thomas-Agnan. *Reproducing Kernel Hilbert Spaces in Probability and Statistics*. Kluwer, 2004.
- K. Fukumizu, A. Gretton, X. Sun, and B. Schölkopf. Kernel measures of conditional dependence. In *Advances in Neural Information Processing Systems 20*, pages 489–496, Cambridge, MA, 2008. MIT Press.
- K. Fukumizu, B. Sriperumbudur, A. Gretton, and B. Schoelkopf. Characteristic kernels on groups and semigroups. In *Advances in Neural Information Processing Systems 21*, pages 473–480, Red Hook, NY, 2009. Curran Associates Inc.
- Kenji Fukumizu, Le Song, and Arthur Gretton. Kernel bayes’ rule. In J. Shawe-Taylor, R.S. Zemel, P. Bartlett, F.C.N. Pereira, and K.Q. Weinberger, editors, *Advances in Neural Information Processing Systems 24*, pages 1737–1745. 2011.
- A. Gretton, O. Bousquet, A.J. Smola, and B. Schölkopf. Measuring statistical dependence with Hilbert-Schmidt norms. In S. Jain, H. U. Simon, and E. Tomita, editors, *Proceedings of the Conference on Algorithmic Learning Theory (ALT)*, pages 63–77. Springer-Verlag, 2005.
- A. Gretton, K. Fukumizu, C.-H. Teo, L. Song, B. Schölkopf, and A. Smola. A kernel statistical test of independence. In *Advances in Neural Information Processing Systems 20*, pages 585–592, Cambridge, MA, 2008. MIT Press.
- A. Gretton, K. Fukumizu, Z. Harchaoui, and B. Sriperumbudur. A fast, consistent kernel two-sample test. In *Advances in Neural Information Processing Systems 22*, Red Hook, NY, 2009a. Curran Associates Inc.
- A. Gretton, K. Fukumizu, and B. Sriperumbudur. Discussion of: Brownian distance covariance. *Ann. Appl. Stat.*, 3(4):1285–1294, 2009b.
- A. Gretton, K. Borgwardt, M. Rasch, B. Schölkopf, and A. Smola. A kernel two-sample test. *J. Mach. Learn. Res.*, 13:723–773, 2012.
- R. Lyons. Distance covariance in metric spaces. arXiv:1106.5758, June 2011.
- A. Müller. Integral probability metrics and their generating classes of functions. *Adv. Appl. Probab.*, 29(2):429–443, 1997.
- M. Reed and B. Simon. *Methods of modern mathematical physics. Vol. 1: Functional Analysis*. Academic Press, San Diego, 1980.
- B. Schölkopf, A. J. Smola, and K.-R. Müller. Kernel principal component analysis. In W. Gerstner, A. Germond, M. Hasler, and J.-D. Nicoud, editors, *ICANN*, volume 1327 of *Lecture Notes in Computer Science*, pages 583–588, Berlin, 1997. Springer-Verlag.
- D. Sejdinovic, A. Gretton, B. Sriperumbudur, and K. Fukumizu. Hypothesis testing using pairwise distances and associated kernels. In *Proceedings of the International Conference on Machine Learning (ICML)*, 2012.
- A. J. Smola, A. Gretton, L. Song, and B. Schölkopf. A Hilbert space embedding for

- distributions. In *Proceedings of the Conference on Algorithmic Learning Theory (ALT)*, volume 4754, pages 13–31. Springer, 2007.
- B. Sriperumbudur. Mixture density estimation via Hilbert space embedding of measures. In *Proceedings of the International Symposium on Information Theory*, pages 1027 – 1030, 2011.
- B. Sriperumbudur, A. Gretton, K. Fukumizu, G. Lanckriet, and B. Schölkopf. Injective Hilbert space embeddings of probability measures. In *Proceedings of the Conference on Learning Theory (COLT)*, pages 111–122, 2008.
- B. Sriperumbudur, K. Fukumizu, A. Gretton, G. Lanckriet, and B. Schoelkopf. Kernel choice and classifiability for RKHS embeddings of probability distributions. In *Advances in Neural Information Processing Systems 22*, Red Hook, NY, 2009. Curran Associates Inc.
- B. Sriperumbudur, A. Gretton, K. Fukumizu, G. Lanckriet, and B. Schölkopf. Hilbert space embeddings and metrics on probability measures. *J. Mach. Learn. Res.*, 11: 1517–1561, 2010.
- B. Sriperumbudur, K. Fukumizu, and G. Lanckriet. Universality, characteristic kernels and rkhs embedding of measures. *J. Mach. Learn. Res.*, 12:2389–2410, 2011.
- I. Steinwart and A. Christmann. *Support Vector Machines*. Springer, 2008.
- G. Székely and M. Rizzo. Testing for equal distributions in high dimension. *InterStat*, (5), November 2004.
- G. Székely and M. Rizzo. A new test for multivariate normality. *J. Multivariate Anal.*, 93:58–80, 2005.
- G. Székely and M. Rizzo. Brownian distance covariance. *Ann. Appl. Stat.*, 4(3):1233–1303, 2009.
- G. Székely, M. Rizzo, and N.K. Bakirov. Measuring and testing dependence by correlation of distances. *Ann. Stat.*, 35(6):2769–2794, 2007.
- K. Zhang, J. Peters, D. Janzing, and B. Schoelkopf. Kernel-based conditional independence test and application in causal discovery. In *Proceedings of the Conference on Uncertainty in Artificial Intelligence (UAI)*, pages 804–813. AUAI Press, Corvallis, OR, USA, 2011.

APPENDIX A: DISTANCE CORRELATION

As described by Székely et al. [2007], the notion of distance covariance extends naturally to that of *distance variance* $\mathcal{V}^2(X) = \mathcal{V}^2(X, X)$ and of *distance correlation* (by analogy with the Pearson product-moment correlation coefficient),

$$\mathcal{R}^2(X, Y) = \begin{cases} \frac{\mathcal{V}^2(X, Y)}{\mathcal{V}(X)\mathcal{V}(Y)}, & \mathcal{V}(X)\mathcal{V}(Y) > 0, \\ 0, & \mathcal{V}(X)\mathcal{V}(Y) = 0. \end{cases}$$

Distance correlation also has a straightforward interpretation in terms of kernels,

$$\begin{aligned}\mathcal{R}^2(X, Y) &= \frac{\mathcal{V}^2(X, Y)}{\mathcal{V}(X)\mathcal{V}(Y)} \\ &= \frac{\gamma_k^2(P_{XY}, P_X P_Y)}{\gamma_k(P_{XX}, P_X P_X) \gamma_k(P_{YY}, P_Y P_Y)} \\ &= \frac{\|\Sigma_{XY}\|_{HS}^2}{\|\Sigma_{XX}\|_{HS} \|\Sigma_{YY}\|_{HS}},\end{aligned}$$

where covariance operator $\Sigma_{XY} : \mathcal{H}_{k_X} \rightarrow \mathcal{H}_{k_Y}$ is a linear operator for which $\langle \Sigma_{XY} f, g \rangle_{\mathcal{H}_{k_Y}} = \mathbb{E}_{XY} [f(X)g(Y)] - \mathbb{E}_X f(X) \mathbb{E}_Y g(Y)$ for all $f \in \mathcal{H}_{k_X}$ and $g \in \mathcal{H}_{k_Y}$, and $\|\cdot\|_{HS}$ denotes the Hilbert-Schmidt norm [Gretton et al. \[2005\]](#). It is clear that $\mathcal{R}$ is invariant to scaling $(X, Y) \mapsto (\epsilon X, \epsilon Y)$, $\epsilon > 0$, whenever the corresponding semimetrics are homogeneous, i.e., whenever $\rho_X(\epsilon x, \epsilon x') = \epsilon \rho_X(x, x')$, and similarly for ρ_Y . Moreover, $\mathcal{R}$ is invariant to translations, $(X, Y) \mapsto (X + x', Y + y')$, $x' \in \mathcal{X}$, $y' \in \mathcal{Y}$, whenever ρ_X and ρ_Y are translation invariant.

APPENDIX B: LINK WITH UNIVERSAL KERNELS

We briefly remark on how our results on equivalent kernels relate to the notion of universal kernels on compact metric spaces in the sense of [Steinwart and Christmann \[2008, Definition 4.52\]](#):

DEFINITION 32. A continuous kernel k on a compact metric space $\mathcal{Z}$ is said to be universal if its RKHS $\mathcal{H}_k$ is dense in the space $C(\mathcal{Z})$ of continuous functions on $\mathcal{Z}$, endowed with the uniform norm.

The family of universal kernels includes the most popular choices in machine learning literature including the Gaussian and the Laplacian kernel. The following characterization of universal kernels is due to [Sriperumbudur et al. \[2011\]](#):

PROPOSITION 33. *Let k be a continuous kernel on a compact metric space $\mathcal{Z}$. Then, k is universal if and only if $\mu_k : \mathcal{M}(\mathcal{Z}) \rightarrow \mathcal{H}_k$ is a vector space monomorphism, i.e.,*

$$\|\mu_k(\nu)\|_{\mathcal{H}_k}^2 = \int \int k(z, z') d\nu(z) d\nu(z') > 0 \quad \forall \nu \in \mathcal{M}(\mathcal{Z}) \setminus \{0\}.$$

As a direct consequence, every universal kernel k is also characteristic, as μ_k is, in particular, injective on the space of probability measures. Now, consider a kernel $\tilde{k}_f$ centered at $f = \mu_k(\nu)$ for some $\nu \in \mathcal{M}(\mathcal{Z})$, such that $\nu(\mathcal{Z}) = 1$. Then $\tilde{k}_f$ is no longer universal, since

$$\begin{aligned}
\left\| \mu_{\tilde{k}_f}(\nu) \right\|_{\mathcal{H}_{\tilde{k}_f}}^2 &= \int \tilde{k}_f(z, z') d\nu(z) d\nu(z') \\
&= \int \int \left[k(z, z') - \int k(w, z) d\nu(w) - \int k(w, z') d\nu(w) \right. \\
&\quad \left. + \int \int k(w, w') d\nu(w) d\nu(w') \right] d\nu(z) d\nu(z') \\
&= (1 - \nu(\mathcal{Z}))^2 \|\mu_k(\nu)\|_{\mathcal{H}_k}^2 \\
&= 0.
\end{aligned}$$

However, $\tilde{k}_f$ is still characteristic, as it is equivalent to k . This means that all kernels of the form (2.12), including the distance kernels, are examples of non-universal characteristic kernels, provided that they generate a semi-metric ρ of strong negative type. In particular, the kernel in (2.10) on a compact $\mathcal{Z} \subset \mathbb{R}^d$ is a characteristic non-universal kernel for $q < 2$. This result is of some interest to the machine learning community, as such kernels have typically been difficult to construct. For example, the two notions are known to be equivalent on the family of translation invariant kernels on $\mathbb{R}^d$ [Sriperumbudur et al., 2011].

ACKNOWLEDGEMENTS

D. Sejdinovic, B. Sriperumbudur and A. Gretton acknowledge support of the Gatsby Charitable Foundation.

GATSBY COMPUTATIONAL NEUROSCIENCE UNIT
 UNIVERSITY COLLEGE LONDON
 17 QUEEN SQUARE, LONDON, WC1N 3AR
 UNITED KINGDOM
 E-MAIL: dino.sejdinovic@gmail.com
bharath@gatsby.ucl.ac.uk
arthur.gretton@gmail.com

THE INSTITUTE OF STATISTICAL MATHEMATICS
 10-3 MIDORICHO, TACHIKAWA
 TOKYO 190-8562
 JAPAN
 E-MAIL: fukumizu@ism.ac.jp